

## Recent ornithological literature from the Caribbean

A regular feature of the Journal of Caribbean Ornithology, this column alerts readers to recent ornithological literature from the Caribbean basin that has appeared elsewhere. We would also like to include any unpublished theses or other reports that may be difficult to find in more universally available abstract services. We invite readers of the Journal of Caribbean Ornithology to alert our compiler, Steven Latta, to other articles that should be highlighted in this section. Our hope is that by providing these summaries we will increase the exchange of knowledge among Caribbean ornithologists and conservationists.

—Steven C. Latta

National Aviary, Allegheny Commons West, Pittsburgh, PA 15212, USA; e-mail: [steven.latta@aviary.org](mailto:steven.latta@aviary.org).

Angelier, F., R.L. Holberton, and P.P. Marra. 2009. Does stress response predict return rate in a migratory bird species? A study of American redstarts and their non-breeding habitat. *Proceedings of the Royal Society B* 276:3545–3551.—The adrenocortical stress response of non-breeding American Redstarts (*Setophaga ruticilla*), wintering in habitats of either high (mangrove) or low suitability (scrub), was measured, and their return rate during the following non-breeding seasons was monitored. Results suggest that in a context-dependent manner, the ability of an individual to physiologically react to stress determines its ability of returning to its non-breeding territory the following winters. E-mail: [fangelier@ucdavis.edu](mailto:fangelier@ucdavis.edu).

Arendt, W. 2011. Hourly laying patterns of the Pearly-eyed Thrasher (*Margarops fuscatus*) in Puerto Rico. *Wilson Journal of Ornithology* 123:624–628.—E-mail: [waynearendt@gmail.com](mailto:waynearendt@gmail.com).

Beason, J.P., C. Gunn, K.M. Potter, R.A. Sparks, and J.W. Fox. 2012. The Northern Black Swift: migration path and wintering area revealed. *Wilson Journal of Ornithology* 124:1–8.—E-mail: [jason.beason@rmbo.org](mailto:jason.beason@rmbo.org).

Beissinger, S.R., J.M. Wunderle, Jr., J.M. Meyers, B.-E. Saether, and S. Engen. 2008. Anatomy of a bottleneck: diagnosing factors limiting population growth in the Puerto Rican Parrot. *Ecological Monographs* 78:185–203.—The role of genetic, demographic, environmental, and catastrophic processes in maintaining the Puerto Rican Parrot (*Amazona vittata*) in a prolonged bottleneck are investigated. Results suggest that four primary factors (reduced hatching success due to inbreeding, failure of adults to nest, nest failure due to non-genetic causes, and reduced survival of adults and juveniles) were responsible for maintaining the bottleneck. E-mail: [jmwunderle@gmail.com](mailto:jmwunderle@gmail.com).

Bibles, B.D., and C.W. Boal. 2012. Morphometric-based sexual determination of Bananaquits (*Coereba flaveola*). *Ornitología Neotropical* 23:507–515.—Morphometric data of the Bananaquit were collected over 8 yrs on Guana Island, British Virgin Islands,

to develop a predictive model based on logistic regression to assign adult Bananaquits to sex. This model classified 96% of validation individuals to the correct sex. E-mail: [clint.boal@ttu.edu](mailto:clint.boal@ttu.edu).

Bouchard, L.C., and M.J. Anderson. 2011. Caribbean Flamingo resting behavior and the influence of weather variables. *Journal of Ornithology* 152:307–312.—Observations of captive American Flamingos (*Phoenicopterus ruber*) sought to investigate the possible functions of unipedal and bipedal resting. Results suggested that flamingos rest on one leg for longer intervals in an attempt to conserve body heat, and that on windy days a bipedal stance is preferred to enhance stability. E-mail: [mandero6@sju.edu](mailto:mandero6@sju.edu).

Brown, A.C., and K. Brindock. 2011. Breeding success and nest site selection by a Caribbean population of Wilson's Plovers. *Wilson Journal of Ornithology* 123:814–819.—E-mail: [abrown@epic-islands.org](mailto:abrown@epic-islands.org).

Brown, D.R., and T.W. Sherry. 2008. Alternative strategies of space use and response to resource change in a wintering migrant songbird. *Behavioral Ecology* 19:1314–1325.—Sedentary and floating behaviors were studied in a wintering population of the Ovenbird (*Seiurus aurocapilla*) in response to manipulated and natural variation in food availability in Jamaica. Results suggest that alternative behaviors represent a trade-off in response to resource availability. E-mail: [david.brown@eku.edu](mailto:david.brown@eku.edu).

Clark, C.J. 2011. Effects of tail length on an escape maneuver of the Red-billed Streamertail. *Journal of Ornithology* 152:397–408.—This study aimed to test whether experimental manipulation of tail length affected the kinematics of a low-speed escape maneuver in the Streamertail (*Trochilus polytmus*) of Jamaica. Experimental observations suggest that the greatly elongated tails of this species may not pose a large cost to low-speed maneuvering flight. E-mail: [christopher.clark@yale.edu](mailto:christopher.clark@yale.edu).

Cooper, N.W., M.A. Thomas, M.B. Garfinkel, K.L. Schneider, and P.P. Marra. 2012. Comparing the precision, accuracy, and efficiency of branch clipping and sweep netting for sampling arthropods in two Jamaican forest types. *Journal of Field Ornithology* 83:381–390.—E-mail: [nathanwands@gmail.com](mailto:nathanwands@gmail.com).

Dalsgaard, B. 2011. Nectar-feeding and pollination by the Cuban Green Woodpecker (*Xiphidiopicus percussus*) in the West Indies. *Ornitología Neotropical* 22:447–451.—E-mail: [b.dalsgaard@zoo.cam.ac.uk](mailto:b.dalsgaard@zoo.cam.ac.uk).

Dalsgaard, B., D.W. Carstensen, A. Kirkconnell, A.M. Martín González, O. Martínez García, A. Timmermann, and W.J. Sutherland. 2012. Floral traits of plants visited by the Bee Hummingbird (*Mellisuga helenae*). *Ornitología Neotropical* 23:143–149.—E-mail: [bo.dalsgaard@bio.ku.dk](mailto:bo.dalsgaard@bio.ku.dk).

Ewert, D.N., K.R. Hall, J.M. Wunderle, Jr., D. Currie, S.M. Rockwell, S.B. Johnson, and J.D. White. 2012. Duration and rate of spring migration of Kirtland's Warblers. *Wilson Journal of Or-*

nithology 124:9–14.—Migration of five Kirtland's Warblers (*Setophaga kirtlandii*) was recorded by observing uniquely color-banded individuals at or near both the beginning and end of spring migration in the Bahamas and Michigan. Average duration of migration was 15.8 days (range 13–23 days) and the average distance traveled was 144.5 km/day (96.1–169.1 km/day). E-mail: dewert@tnc.org.

Gilardi, J. 2011. Raise...release...repeat. *PsittaScene* 23:3–5.—An accounting of the efforts to conserve the Puerto Rican Parrot, including captive breeding at the Luquillo Aviary in the El Yunque Rainforest, and at the Vivaldi (Rio Abajo) Aviary in the Rio Abajo Forest. As of the start of 2011 the wild population of parrots was 34–40 individuals in the Rio Abajo State Forest, and 20–30 in the Caribbean National Forest. The captive population is now 269 parrots, split evenly between both aviaries. E-mail: unavailable.

Gowda, V., E.J. Temeles, and W.J. Kress. 2012. Territorial fidelity to nectar sources by Purple-throated Caribs, *Eulampis jugularis*. *Wilson Journal of Ornithology* 124:81–86.—First record of territorial site-fidelity across multiple years by Purple-throated Caribs on St. Kitts, Dominica, and St. Vincent. Males remained near patches of their main nectar resources even when they were not in flower. E-mail: vinitagowda@gmail.com.

Jirinec, V., B.R. Campos, and M.D. Johnson. 2011. Roosting behaviour of a migratory songbird on Jamaican coffee farms: landscape composition may affect delivery of an ecosystem service. *Bird Conservation International* 21:353–361.—Nocturnal tracking revealed Black-throated Blue Warblers (*Setophaga caeruleascens*) moved outside diurnal foraging ranges on the farms to roost in forested habitat patches, sometimes up to 1 km away. These findings reflect the ecological connectivity between coffee habitats and the adjacent landscape. E-mail: vj2@humboldt.edu.

Juhant, M.A. 2012. Where to watch raptor migration in the Caribbean. *Neotropical Birding* 11:4–15.—E-mail: matias\_juhant@yahoo.com.ar.

Kennedy, C.M., E.H. Campbell Grant, M.C. Neel, W.F. Fagan, and P.P. Marra. 2011. Landscape matrix mediates occupancy dynamics of Neotropical avian insectivores. *Ecological Applications* 21:1837–1850.—Multi-season occupancy models were used to determine the relative influence of patch area, patch isolation, within-patch vegetation structure, and landscape matrix on occupancy dynamics of nine Neotropical insectivorous birds in 99 forest patches embedded in four matrix types (agriculture, suburban development, bauxite mining, and forest) in central Jamaica. Within-patch vegetation structure and the matrix type between patches were more important than patch area and patch isolation in determining local colonization and local extinction probabilities, and the effects of patch area, isolation, and vegetation structure on occupancy dynamics tended to be matrix and species dependent. E-mail: cmk6@umd.edu.

Kirwan, G.M. 2012. Jamaica - "the fairest island." *Neotropical Birding* 11:47–55.—E-mail: GMKirwan@aol.com.

LaPergola, J.B., J.L. Mortensen, and R.L. Curry. 2011. Nest, eggs, and nesting behavior of the Gray Trembler (*Cinlocerthia gutturalis*) on St. Lucia, West Indies. *Wilson Journal of Ornithology* 123:390–395.—E-mail: jbl96@cornell.edu.

Latta, S.C. 2012. Avian research in the Caribbean: past contributions and current priorities. *Journal of Field Ornithology*

83:107–121.—E-mail: steven.latta@aviary.org.

Lefevre, K.L., S. Sharma, and F.H. Rodd. 2012. Moderate human disturbance of rain forest alters composition of fruiting plant and bird communities. *Biotropica* 44:427–436.—On Tobago, seemingly moderate human disturbance has led to substantial changes in the plant and bird assemblages of the rainforest, underscoring the impact of even moderate activity on community composition. E-mail: k.lefevre@alumni.utoronto.ca.

Longrich, N.R., and S.L. Olson. 2011. The bizarre wing of the Jamaican flightless ibis *Xenicibis xymptithecus*: a unique vertebrate adaptation. *Proceedings of the Royal Society B* 278:2333–2337.—Fossil evidence is presented showing a radically different metacarpus that is elongated and grossly inflated with extremely thick walls. The authors suggest that this bizarre morphology of the wing was used as a club, perhaps in intraspecific battles. E-mail: nicholas.longrich@yale.edu.

Mathys, B.A. 2011. First record of Aplomado Falcon (*Falco femoralis*) for the West Indies. *Wilson Journal of Ornithology* 123:179–180.—E-mail: Blake.Mathys@stockton.edu.

Monceau, K., R. Wattier, F.-X. Dechaume-Moncharmont, S. Motreuil, and F. Cézilly. 2011. Territoriality versus flocking in the Zenaida Dove (*Zenaida aurita*): resource polymorphism revisited using morphological and genetic analyses. *Auk* 128:15–25.—The term resource polymorphism refers to the existence of alternative phenotypes in relation to resource use. On Barbados, although Zenaida Doves usually defend year-round territories, birds can also be observed foraging at seed-storage sites in large flocks with little aggression. It has been suggested that this represents a case of resource polymorphism driven by competition for territories. This paper uses new data to revisit the evidence for resource polymorphism in Zenaida Doves on Barbados, but overall the existence of resource polymorphism is questioned. E-mail: karine.monceau@u-bourgogne.fr.

Mones Espin, R., and L. Garcia Rivera. 2010. First breeding record of Black Swift *Cypseloides niger* in Cuba. *Cotinga* 32:146–147.—E-mail: montesninin@ups.perla.inf.cu.

Pérez Mena, E.E., and E.C. Mora. 2011. Geographic song variation in the non-oscine Cuban Tody (*Todus multicolor*). *Wilson Journal of Ornithology* 123:76–84.—Cuban Todies emitted three types of sounds across the Cuban archipelago. Evidence of geographic variation was found with todies grouped into two main clusters corresponding to western and eastern Cuba. This geographic song variation may indicate genetic differences in this sedentary forest bird, and the existence of two "incipient species" of todies in Cuba. E-mail: jr Rubio@infomed.sld.cu.

Price, M.R., V.A. Lee, and W.K. Hayes. 2011. Population status, habitat dependence, and reproductive ecology of Bahama Orioles: a critically endangered synanthropic species. *Journal of Field Ornithology* 82:366–378.—In light of the recent elevation of the critically endangered Bahama Oriole (*Icterus northropi*) to species status, this evaluates their population status, habitat use, and breeding ecology. E-mail: mrprice@llu.edu.

Reudink, M.W., P.P. Marra, T.K. Kyser, P.T. Boag, K.M. Langin, and L.M. Ratcliffe. 2009. Non-breeding season events influence sexual selection in a long-distance migratory bird. *Proceedings of the Royal Society B* 276:1619–1626.—This study examines whether carry-over effects from the non-breeding season can influence the process of sexual selection in the migratory Amer-

ican Redstart. Partially based on data from Jamaica, results suggest that the process of sexual selection may be influenced by events interacting throughout the annual cycle. E-mail: mattreudink@gmail.com.

Ricklefs, R.E., and E. Bermingham. 2007. The causes of evolutionary radiations in archipelagoes: passerine birds in the Lesser Antilles. *American Naturalist* 169:285–297.—Passerine avifaunas of the Hawaiian and Galapagos archipelagoes, which have supported well-known radiations of birds, are compared with those of the Lesser Antilles, which have not. Analysis of genetic divergence among island populations in the Lesser Antilles reveals evidence of both prolonged independent evolution and re-expansion of differentiated island populations through the archipelago but little evidence of secondary sympatry of divergent genetic lineages. E-mail: ricklefs@umsl.edu.

Rockwell, S.M., C.I. Bocetti, and P.P. Marra. 2012. Carry-over effects of winter climate on spring arrival date and reproductive success in an endangered migratory bird, Kirtland's Warbler (*Setophaga kirtlandii*). *Auk* 129:744–752.—Males arrive on breeding grounds later following drier winters, and first-time breeders are more sensitive to changes in rainfall on the wintering grounds than experienced adults. Regardless of age, drier winters and delayed arrival and nest initiation are significantly associated with fewer offspring fledged. E-mail: rockwell@umd.edu.

Sly, N.D., A.K. Townsend, C.C. Rimmer, J.M. Townsend, S.C. Latta, and I.J. Lovette. 2011. Ancient islands and modern invasions: disparate phylogeographic histories among Hispaniola's endemic birds. *Molecular Ecology* 20:5012–5024.—Multilocus comparative phylogeography techniques are used to examine the pattern and history of divergence in 11 endemic birds representing potential within-island speciation events. Phylogeographic divergence within or between closely related species was correlated with the likely distribution of ancient sea barriers that once divided Hispaniola into several smaller paleo-islands, and no evidence was found that ecological or topographical complexity generated diversity, either by creating open niches or by restricting long-term gene flow. E-mail: nds22@cornell.edu.

Smith, J.A.M., L.R. Reitsma, and P.P. Marra. 2011. Influence of moisture and food supply on the movement dynamics of a non-breeding migratory bird (*Parkesia noveboracensis*) in a seasonal landscape. *Auk* 128:43–52.—Northern Waterthrushes (*Parkesia noveboracensis*) were radio-tracked in four habitats in Puerto Rico during two winters to determine the ecological determinants of diurnal space use and overwinter site-persistence. The primary determinants of movement probability were moisture and food availability on each bird's home range. E-mail: joseph\_smith@tnc.org.

Smith, J.A.M., L.R. Reitsma, and P.P. Marra. 2011. Multiple space-use strategies and their divergent consequences in a nonbreeding migratory bird (*Parkesia noveboracensis*). *Auk* 128: 53–60.—Relationships among space-use patterns, home-range attributes, and individual characteristics of Northern Waterthrush were investigated to determine the consequences of different space-use strategies for overwinter physical condition. Intraspecific competition played an important role in determining where individuals initially and eventually settled, and territoriality, characterized by aggression, site-persistence, and exclusive

home ranges, often resulted in benefits, especially for males. E-mail: joseph\_smith@tnc.org.

Studds, C.E., T.K. Kyser, and P.P. Marra. 2008. Natal dispersal driven by environmental conditions interacting across the annual cycle of a migratory songbird. *Proceedings of the National Academy of Science* 105:2929–2933.—Using stable-hydrogen isotope ratios in feathers of American Redstarts captured as immature birds and again as adults, habitat use during the first tropical non-breeding season is shown to interact with latitudinal gradients in spring phenology on the temperate breeding grounds to influence the distance traveled on the initial spring migration and the direction of natal dispersal. In contrast, adult redstarts showed considerable site fidelity between breeding seasons, indicating that environmental conditions did not affect dispersal patterns after the first breeding attempt. E-mail: marrap@si.edu.

Toms, J.D., J. Faaborg, and W.J. Arendt. 2012. Climate change and birds in the forgotten tropics: the importance of tropical dry forests. *Ibis* 154:632–634.—E-mail: judith.toms@zoho.com.

Townsend, J.M., C.C. Rimmer, A.K. Townsend, and K.P. McFarland. 2011. Sex and age ratios of Bicknell's Thrush wintering in Hispaniola. *Wilson Journal of Ornithology* 123:367–372.—The proportion of male Bicknell's Thrush (*Catharus bicknelli*) on Hispaniola was 0.64, which is comparable to the known male bias in breeding areas. The proportion of males varied geographically on Hispaniola, suggesting some level of habitat segregation, with males preferentially occupying cloud forest sites characterized by a thick understory of vines and saplings.—E-mail: townsend.jason.m@gmail.com.

Townsend, J.M., C.C. Rimmer, and K.P. McFarland. 2012. Radio-transmitters do not affect seasonal mass change or annual survival of wintering Bicknell's Thrushes. *Journal of Field Ornithology* 83:295–301.—Studies of transmitters on Bicknell's Thrushes wintering in the Dominican Republic indicate that attaching light-weight transmitters (< 5% of body mass) to wintering Bicknell's Thrushes did not adversely affect either body condition or annual survival. E-mail: townsend.jason.m@gmail.com.

Townsend, J.M., C.C. Rimmer, K.P. McFarland, and J.E. Goetz. 2012. Site-specific variation in food resources, sex ratios, and body condition of an overwintering migrant songbird. *Auk* 129:683–690.—Available food resources, sex ratios, and body condition of territorial Bicknell's Thrushes wintering in the Dominican Republic were studied at two ecologically distinct wet-forest sites. Results suggest that there are sex-specific advantages of wintering in these two habitats and that both are critical to supporting the full demographic structure of Bicknell's Thrush populations. E-mail: townsend.jason.m@gmail.com.

Villard, P., P. Feldmann, A. Ferchal, and C. Pavis. 2010. Population size and habitat associations of the endemic Guadeloupe Woodpecker. *Journal of Field Ornithology* 81:278–286.—E-mail: pascalvillard@yahoo.fr.

Vilella, F.J., and W.F. Nimitz. 2012. Spatial dynamics of the Red-tailed Hawk in the Luquillo Mountains of Puerto Rico. *Wilson Journal of Ornithology* 124:758–766.—Movements and resource use of 21 radio-marked Red-tailed Hawks (*Buteo jamaicensis*) were studied by delineating home range, core area shifts, and macrohabitat use in northeastern Puerto Rico. Find-

ings suggest fragmentation of contiguous forest may benefit the species. E-mail: fvilella@cfr.msstate.edu.

White, R.L., T.J. Baptiste, A. Dornelly, M.N. Morton, M.J. O'Connell, and R.P. Young. 2012. Population responses of the Endangered White-breasted Thrasher *Ramphocinclus brachyurus* to a tourist development in Saint Lucia – conservation implications from a spatial modelling approach. *Bird Conservation International* 22:468–485.—A four year dataset (2006–2009) and landscape-scale environmental variables were used to: a) identify, characterize and map spatio-temporal patterns of thrasher encounter rates, b) determine landscape-scale environmental variables that influence such patterns, and c) produce an island-wide predictive map of potentially suitable habitat. Findings suggest that it is vital that patches of suitable dry forest adjacent to the tourist development are protected and contigu-

ous natural habitat inside the tourist development is preserved. E-mail: rlw22@kent.ac.uk.

Wiancko, E., E. Nol, A. Parada, and D.M. Burke. 2011. Landbird richness and abundance in three coastal habitats near resorts in Cayo Coco, Cuba. *Condor* 113:41–51.—E-mail: ewiancko@yahoo.ca.

Young, R.P., T.J. Baptiste, A. Dornelly, H. Temple, H. Whitehead, H.G. Young, and M.N. Morton. 2010. Potential impacts of tourist developments in St Lucia on the Endangered White-breasted Thrasher *Ramphocinclus brachyurus*. *Bird Conservation International* 20:354–364.—E-mail: matthew.morton@durrell.org.

Zuccon, D. 2011. A new name for the Montserrat Forest Thrush. *Bulletin of the British Ornithologists' Club* 131:199–200.—E-mail: dario.zuccon@libero.it.