

RECENT ORNITHOLOGICAL LITERATURE

Readers are invited to submit literature citations that should be highlighted in this section to STEVEN C. LATTA, *National Aviary, Allegheny Commons West, Pittsburgh, PA 15212, USA*; e-mail: steven.latta@aviary.org.

ACOSTA, M., L. MUGICA, D. BLANCO, B. LOPEZ-LANUS, R. ANTUNES DIAS, L. W. DOODNATH, AND J. HURTADO. 2010. Birds of rice fields in the Americas. *Waterbirds* 33:105–122.—Data on waterbird and landbird use of rice paddies was compiled from Argentina, Brazil, Colombia, Costa Rica, Cuba, Surinam, Trinidad and Tobago, Uruguay, USA, and Venezuela. At least 169 waterbirds and 166 landbirds have been recorded in the paddies of the region. In all countries, rice fields are considered important feeding areas and heavily used as migratory stop-over and wintering sites. E-mail: macosta@fbio.uh.cu.

BELTRAN, J. W., J. M. WUNDERLE, AND W. J. ARENDT. 2010. Changes in home range of breeding and post-breeding male Pearly-eyed Thrashers in the Luquillo Mountains of Puerto Rico. *Ornitología Neotropical* 21:409–423.—E-mail: williambeltran@gmail.com.

DEXTER, K. G. 2010. The influence of dispersal on macroecological patterns of Lesser Antillean birds. *Journal of Biogeography* 37:2137–2147.—Dispersal is often assumed to be a major force in shaping macroecological patterns. Here macroecological patterns of distribution for Lesser Antillean birds of the rain forest and dry forest are described, and then population genetic data used to assess if differences in dispersal ability could be responsible for the groups' contrasting patterns. Results suggest that differences in dispersal ability are a plausible explanation for the contrasting macroecological patterns. But historical factors, such as the taxon cycle and Pleistocene climate fluctuations, may have also played a role in shaping the distribution patterns. E-mail: kgdexter@gmail.com.

FAABORG, J., R. T. HOLMES, A. D. ANDERS, K. L. BILDSTEIN, K. M. DUGGER, S. A. GAUTHREAUX, P. HEGLUND, K. A. HOBSON, A. E. JAHN, D. H. JOHNSON, S. C. LATTA, D. J. LEVEY, P. P. MARRA, C. L. MERKORD, E. NOL, S. I. ROTHSTEIN, T. W. SHERRY, T. S. SILLETT, F. R. THOMPSON, AND N. WARNOCK. 2010. Recent advances in understanding migration systems of New World land birds. *Ecological Monographs* 80:1–48.—E-mail: [faaborgj@](mailto:faaborgj@missouri.edu)

missouri.edu.

FAABORG, J., R. T. HOLMES, A. D. ANDERS, K. L. BILDSTEIN, K. M. DUGGER, S. A. GAUTHREAUX JR., P. HEGLUND, K. A. HOBSON, A. E. JAHN, D. H. JOHNSON, S. C. LATTA, D. J. LEVEY, P. P. MARRA, C. L. MERKORD, E. NOL, S. I. ROTHSTEIN, T. W. SHERRY, T. S. SILLETT, F. R. THOMPSON III, AND N. WARNOCK. 2010. Managing migratory landbirds in the New World: do we know enough? *Ecological Applications* 20:398–418.—E-mail: faaborgj@missouri.edu.

GALVEZ AGUILERA, X., AND F. CHAVEZ-RAMIREZ. 2010. Distribution, abundance, and status of Cuban Sandhill Cranes (*Grus canadensis nesiototes*). *Wilson Journal of Ornithology* 122:556–562.—E-mail: fchavez@whoopingcrane.org.

HAYES, F. E., B. SANASIE, AND I. SAMAD. 2009. Status and conservation of the critically endangered Trinidad Piping-Guan *Aburria pipile*. *Endangered Species Research* 7:77–84.—E-mail: fhayes@puc.edu.

HAYES, F. E., C. L. SHAMEERUDEEN, B. SANASIE, B. D. HAYES, C. L. RAMJOHN, AND F. B. LUCAS. 2009. Ecology and behaviour of the critically endangered Trinidad Piping-Guan *Aburria pipile*. *Endangered Species Research* 6:223–229.—E-mail: fhayes@puc.edu.

HAYES, W. K., E. D. BRACEY, M. R. PRICE, V. ROBINETTE, E. GREN, AND C. STAHALA. 2010. Population status of Chuck-will's-widow (*Caprimulgus carolinensis*) in the Bahamas. *Wilson Journal of Ornithology* 122:381–384.—E-mail: whayes@llu.edu.

KENNEDY, C. M., P. P. MARRA, W. F. FAGAN, AND M. C. NEEL. 2010. Landscape matrix and species traits mediate responses of Neotropical resident birds to forest fragmentation in Jamaica. *Ecological Monographs* 80:651–669.—The authors investigated whether Neotropical resident bird communities in limestone forest patches differed if they were embedded in three different human-dominated matrix types (agriculture, peri-urban development, and bauxite mining) relative to sites in continuous forest in central Jamaica. Species richness, community composition, and abundances were matrix-dependent, with agricultural landscapes supporting greater avian diversity and more intact community assemblages than either peri-urban or bauxite landscapes. This study reinforces the importance of differentiating among land cover and land uses in frag-

mentation research and lends support to the hypothesis that resource availability may be a primary factor driving Neotropical bird responses to fragmentation. E-mail: cmk6@umd.edu.

LATTA, S. C., AND R. E. RICKLEFS. 2010. Prevalence patterns of avian Haemosporidia on Hispaniola. *Journal of Avian Biology* 41:25–33.—E-mail: steven.latta@aviary.org.

NIJMAN, V. 2010. The importance of small wetlands for the conservation of the endemic Caribbean Coot, *Fulica caribaea*. *Caribbean Journal of Science* 46:112–115.—This study used data from 19 sites in the Netherlands Antilles, and records of 49 sites > 5 ha throughout its range, to analyze the relationship between number of coots and wetland size, and the relationships between wetland size, inclusion in the protected area network, and threats to the sites or coots.—E-mail: vnijman@brookes.ac.uk.

MARIN E., G., Y. CARVAJAL M., AND J. MUÑOZ G. 2010. Primer registro de *Thalassarche chlororhynchos* en la cuenca del Mar Caribe, Cotinga 32:159–160.—E-mail: gediom@yahoo.com.

PAULINO, M. M., D. A. MEJIA, AND S. C. LATTA. 2010. A new review of the status of the Caribbean Flamingo (*Phoenicopterus ruber*) in the Dominican Republic and Haiti. *Flamingo* 18:62–66.—E-mail: steven.latta@aviary.org.

RODRIGUEZ-SANTANA, F. 2010. Reports of Cooper's Hawks (*Accipiter cooperii*), Swainson's Hawks (*Buteo swainsoni*), and Short-tailed Hawks (*Buteo brachyurus*) in Cuba. *Journal of Raptor Research* 44:146–150.—E-mail: freddy@bioeco.ciges.inf.cu.

RUSELLO, M. A., C. STAHALA, D. LALONDE, K. L. SCHMIDT, AND G. AMTO. 2010. Cryptic diversity and conservation units in the Bahama Parrot. *Conservation Genetics* 11:1809–1821.—Mitochondrial DNA-based population genetic and phylogenetic analyses revealed the distinctiveness of the Abaco, Inagua, and now extirpated Acklins populations, detecting diagnostic character support and reciprocal monophyly indicative of three phylogenetic spe-

cies. Congruent results were obtained for the Abaco and Inagua populations based on Bayesian clustering analyses of microsatellite genotypic data. Overall, results suggest that the conservation status of the Abaco phylogenetic species should be immediately elevated to reflect its historical isolation, recent population decline, and continued threats to its persistence. E-mail: michael.russello@ubc.ca.

SLY, N. D., A. K. TOWNSEND, C. C. RIMMER, J. M. TOWNSEND, S. C. LATTA, AND I. J. LOVETTE. 2010. Phylogeography and conservation of the endemic Hispaniolan Palm-Tanagers (Aves: *Phaenicophilus*). *Conservation Genetics* 11:2121–2129.—A multilocus phylogeographic approach was used to identify evolutionarily distinct populations of *Phaenicophilus poliocephalus* and *P. palmarum* on Hispaniola. Results support the recognition of two palm-tanager species, confirming *P. poliocephalus* as Haiti's only endemic bird species and underscoring the need to protect the species's single primary forest reserve. E-mail: nds22@cornell.edu.

TEMELES, E. J., C. R. KOULOURIS, S. E. SANDER, AND W. J. KRESS. 2009. Effect of flower shape and size on foraging performance and trade-offs in a tropical hummingbird. *Ecology* 90:1147–1161.—E-mail: ejtemeles@am-herst.edu.

TOWNSEND, J. M., C. C. RIMMER, AND K. P. MCFARLAND. 2010. Winter territoriality and spatial behavior of Bicknell's Thrush (*Catharus bicknelli*) at two ecologically distinct sites in the Dominican Republic. *Auk* 127:514–522.—Radiotelemetry was used to investigate the spatial behavior of wintering Bicknell's Thrushes at a mid-elevation rainforest site and a high-elevation cloud-forest site. Blood stable carbon isotopes and fecal samples were analyzed to compare thrush diets at these sites. Exclusive territoriality was the predominant winter social system, and we suggest that both arthropods and fruit are defensible resources for wintering Bicknell's Thrushes. E-mail: jatownse@syr.edu.

REVIEWERS OF VOLUME 26

We thank the following individuals for reviewing manuscripts (more than one indicated with a numeral) for volume 26: Martin Acosta (5), Matthew J. Anderson, Nicolas Barré, Adam C. Brown, P. A. Buckley (2), Charles T. Collins, Dennis Denis (2), Andrew Dobson (2), Steven D. Emslie, Cyril Eraud,

David Ewert, Hiram González, Ruud van Halewijn (2), William K. Hayes, Ariam Jiménez (3), John Lloyd, Douglas B. McNair, Steven G. Mlodinow, Lourdes Mugica, William L. Murphy, David W. Steadman, and Jason M. Townsend.